

80 mm

140 mm

Hyalin Gel

(Hyaluronic Acid Sodium 0.2%)

ہائیلن جیل
(ہائیلورونک ایسڈ سوڈیم ۰.۲%)

Composition:

Each gram contains:

Hyaluronic Acid Sodium Salt..... 2mg

CLINICAL PHARMACOLOGY

Mechanism of Action:

The principal component is the sodium salt of hyaluronic acid (0.2%). The sodium hyaluronate (Hyalastine) is derived from a natural fermentation process. Hyaluronic acid is a biological polysaccharide (glycosaminoglycan) and is a major component of the extracellular matrix of the connective tissue. Due to its hydrophilic properties, it provides a hydrated environment around and between cells, thus facilitating cell migration

INDICATIONS:

Hyalin gel is indicated for the treatment of skin irritations and lesions. Specifically, it is intended for the coverage of acute and chronic wounds (abrasions, donor and postoperative incisions, first and second degree burns, vascular and metabolic ulcers and bedsores) and ensures a hydrated environment in the wound, protecting it from abrasions, friction and dryness.

DOSAGE & ADMINISTRATION

The wounds or ulcers should be cleaned and disinfected before treatment. Apply a thin layer of gel to the wound surface, two or three times a day for no more than 30 consecutive days. The treated area should be covered with sterile gauze and a sterile bandage.

ROUTE OF ADMINISTRATION

For topical use only.

ADVERSE REACTIONS

The following local adverse reactions may occasionally occur: rash at the application site, itching, burning sensation, skin irritation, dryness, erythema, dermatitis, skin discoloration. The symptoms are usually transient, mild in severity and rarely require early discontinuation of treatment.

CONTRAINDICATIONS

Do not administer to patients with known hypersensitivity to this product.

WARNINGS AND PRECAUTIONS

Avoid direct contact of container with the affected area.

Each container of Hyalin should be used by one patient only in order to reduce the risk of cross infection.

Discard 2 months after first opening.

In case of pregnancy, breastfeeding and use in children, consult your doctor before using the product.

INTERACTIONS

Do not use concomitantly with disinfectants containing quaternary ammonium salts because hyaluronic acid can precipitate in their presence.

The concomitant topical treatment of wounds with antibiotics or other local agents has never given rise to interactions or incompatibilities with Hyalin.

OVERDOSAGE

If conditions worsen, consult your physician immediately. Keep this product out of the reach of children. The prolonged use of the product may give rise to sensitization phenomena. Should this happen, discontinue the treatment and follow a suitable therapy. Do not use the product after expiration date reported on the package.

PRESENTATION

Hyalin gel is available in aluminum tube of 30g.

Dosage and Direction for use: Apply a thin layer of gel to the wound surface for two or three times a day for not more than 30 consecutive days or as directed by the physician.

Instructions:

- Store below 30° C in a dry place, protect from light.
- To be dispensed on the prescription of a registered medical practitioner only.
- Keep out of the reach of children

خوارک اور استعمال کا طریقہ:

جیل کی چمکی تھوڑی سی مقدار چمکی میں ۳ بار لگائیں۔ لگاتار ۳۰ دن سے زیادہ استعمال نہ کریں۔ ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

ہدایات:

دوا کو ۳۰ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر روشنی سے اور نم ہونے سے بچا کر خشک جگہ پر رکھیں۔ نشتے سے پرہیز کریں۔

بچوں کی پہنچ سے دور رکھیں۔ صرف رجسٹرڈ ڈاکٹر کے نسخے پر ہی فروخت کریں۔

Manufactured by:

Kaizen Pharmaceuticals (Pvt.) Ltd.

E-127-129, North Western Industrial Zone,
Bin Qasim, Karachi-75020, Pakistan.

Art No. 1726