Profylline

(Doxofylline)

پروفائلین

(ڈوکسوفائلین)

400mg Tablets &
100mg / 5ml Syrup

Each tablet contains:
Doxofylline (Mfg. Specs.) ….. 400 mg
Product Complies Innovator's Specs.

Each 5 ml contains:
Doxofylline (Mfg. Specs.) ….. 100 mg
Product Complies Innovator's Specs.

Indications:
Asthma, Bronchospasm, chronic osbstructive pulmonary disease and complicated cough.

Dosage:
Tablets: Elderly Patients: 1/2 tablet two - three times daily.
Adults: 1 tablet two - three times daily or as prescribed by the physician.
Syrup: Elderly: 10 ml (2 teaspoonful), two - three times daily.
Children greater than 12 years of age: 10 ml (2 teaspoonful), two - three times daily.
Children less than 12 years of age: 6 – 9 mg/kg twice daily or as directed by the physician.

Contraindications:
This product is contraindicated in individuals who have shown hypersensitivity to its components. It is also contraindicated in patients with acute myocardial infarction, hypotension and in lactating women.

Undesirable Effects:
After xanthine administration, nausea, vomiting, epigastric pain, cephalalgia, irritability, insomnia, tachycardia, extrasystole, tachypnea, and occasionally hyperglycemia and albuminuria, may occur, if a potential oral overdose is established, the patient may present with severe arrhythmias and seizure; these symptoms could be the first sign of an intoxication. Adverse reactions may occur from the withdrawl of treatment; a lower dose rechallenge may start only after the advice of physician.

Special Precautions:
The half-life of xanthine derivatives is influenced by a number of known variables. It may be prolonged in patients with liver disease. In patients with congestive heart failure, in those affected with chronic obstructive lung disease or concomitant infection, and in those patients taking certain other drugs (erythromycin, troleandomycin, lincomycin, and other antibiotics of the same group, allopurinol, cimetidine, propranolol, and anti-flu vaccine). In these cases, a lower dose of Doxofylline may be needed. Phenytoin, other anti-convulsants and smoking may cause an increase in clearance with a shorter mean half-life: in these cases higher doses of Doxofylline may be needed. Use with cautions in patients with hypoxemia, hyperthyroidism, liver disease, renal disease, in those with history of peptic ulcer and in elderly.
Frequently, patients with congestive heart failure have markedly prolonged drug serum levels following discontinuation of the drug.

Use in Pregnancy and Lactation:
Animal reproduction studies indicate that Doxofylline does not cause fetal harm when administered to pregnant animals nor can affect reproduction capacity, However, since there is limited experience in humans during pregnancy, xanthines should be given to a pregnant woman only if clearly needed. Doxofylline is contraindicated in nursing mothers.

Drug Interaction and Others:
Doxofylline should not be administered together with other xanthine derivatives, including beverages and foods containing caffeine. Toxic synergism with ephedrine has been documented for xanthines. Concomitant therapy with erythromycin, troleandomycin, lincomycin, clindamycin, allopurinol, cimetidine, propranolol and anti-flu vaccine may decrease the hepatic clearance of xanthines causing an increase in blood levels.

Presentation:
Profylline 400 mg tablets in blister pack of 10's & 20's.
Profylline 100mg / 5 ml syrup in 60 ml and 120 ml bottle.

* Detail information is available on request.

Dosage as directed by physician. To be dispensed on the prescription of a registered medical practitioner only. Store below 30°C in a dry place, protect from light. Keep out of the reach of children. Keep bottle tightly closed.

- **خوراک:** ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔
- دوا کو ۳۰ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر روشنی سے بچا کر خشک جگہ پر رکھیں۔
- صرف رجسٹرڈ ڈاکٹر کے نسخے پر ہی فروخت کریں۔
- بچوں کی پہنچ سے دور رکھیں۔
- ڈھکن کو اچھی طرح بند رکھیں۔

Manufactured by:
Kaizen Pharmaceuticals (Pvt.) Ltd.
E-127-129, North Western Industrial Zone,
Bin Qasim, Karachi-75020, Pakistan.

ART No. 009/3