· If you have a high blood potassium level, renal disorder, liver disorder or cerebrovascular disorder.
· If you are a preoperative patient, receiving hemodialysis or on a regimen of low-salt diet.
· If you are pregnant, possibly pregnant or breastfeeding.
· If you are taking any other medicinal products. (Some medicines may interact
to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES
· Dizziness and light-headedness may occur due to an excessive decrease in blood pressure. Therefore, pay close attention when working at heights or operating dangerous machinery such as driving a car.

POSSIBLE ADVERSE REACTIONS
The most commonly reported adverse reactions include rash, eczema, itching, dizziness, headache and diarrhea. If any of these symptoms occur, consult with your doctor or pharmacist.
The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.
· Facial/tongue/throat swelling, breathlessness [angioedema]
· Feeling cold, vomiting, falling unconscious [shock, syncope, loss of consciousness]
· Decreased urination output, edema of face and hands/feet, fever [acute renal disorder]
· Numbness of hands/feet and lips, muscle weakness [hyperkalemia]
· General malaise, loss of appetite, yellowing of the skin/white of the eyes [liver dysfunction]
· Muscle pain, lassitude, reddish brown urine [rhabdomyolysis]
The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

PRESENTATION
Myazil 10mg is available in blister packs of 20's.
Myazil 20mg is available in blister packs of 30's.
Myazil 40mg is available in blister packs of 20's.

STORAGE CONDITIONS AND INSTRUCTIONS
· Store below 30°C in a dry place, protect from light.
· To be dispensed on the prescription of a registered medical practitioner only.
· Keep out of the reach of children.

ہدایات:
- خوراک ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔
- دوا کو ۳۰ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر روشنی سے بچا کر خشک جگہ پر رکھیں۔
- صرف رجسٹرڈ ڈاکٹر کے نسخے پر ہی فروخت کریں۔
- بچوں کی پہنچ سے دور رکھیں۔

Manufactured by:
Kaizen Pharmaceuticals (Pvt.) Ltd.
E-127-129, North Western Industrial Zone,
Bin Qasim, Karachi-75020, Pakistan.

Art no: 841/1

Myazil

(Azilsartan)

مائی ایزل

(ایزل سارٹن)

10mg, 20mg and 40mg Tablets

۱۰ ملی گرام ۲۰ ملی گرام اور
۴۰ ملی گرام گولیاں

Composition:
Myazil tablet 10mg
Each film coated tablet contains:
Azilsartan 10mg

Myazil tablet 20mg
Each film coated tablet contains:
Azilsartan 20mg

Myazil tablet 40mg
Each film coated tablet contains:
Azilsartan 40mg

WARNING: FETAL TOXICITY
Myazil can cause serious (possibly fatal) harm to an unborn baby if used during pregnancy. It is important to prevent pregnancy while taking this medication. Consult your doctor for more details and to discuss the use of reliable forms of birth control while taking this medication If you are planning pregnancy, become pregnant, or think you may be pregnant, contact your doctor right away. Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus.

INDICATION
Myazil is indicated for the treatment of high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Myazil belongs to a class of drugs called angiotensin receptor blockers (ARBs). It works by relaxing blood vessels so that blood can flow more easily.

MECHANISM OF ACTION
This medicine lowers blood pressure by reducing peripheral vascular resistance via antagonizing receptor of angiotensin II, an intrinsic substance which raises blood pressure by constricting blood vessels.

POSOLOGY AND METHOD OF ADMINISTRATION
· **Your dosing schedule prescribed by your doctor is (to be written by a healthcare professional)**
· In general, for adults, take 20 mg of the active ingredient at a time, once a day. The dosage may be adjusted according to the age or symptoms, but should not exceed 40 mg a day. This preparation contains 10 mg of the active ingredient in a tablet. Strictly follow the instructions.
· If you miss a dose, take the missed dose as soon as possible. However, wait at least 8 hours to take the next dose. You should never take two doses at one time.
· If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
· Do not stop taking this medicine unless your doctor instructs you to do so.

SPECIAL PRECAUTIONS
The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.
· If you have previously experienced any allergic reactions (itching, rash, etc.) to any medicines or foods.